

## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents  
United States Patent and Trademark  
Office  
Box PCT  
Washington, D.C.20231  
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

<b>Date of mailing</b> (day/month/year) 15 September 2000 (15.09.00)	
<b>International application No.</b> PCT/US99/29213	<b>Applicant's or agent's file reference</b> 9706-0004PC
<b>International filing date</b> (day/month/year) 09 December 1999 (09.12.99)	<b>Priority date</b> (day/month/year) 31 December 1998 (31.12.98)
<b>Applicant</b> DUAN, Lingxun	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

14 July 2000 (14.07.00)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer Ingrid Aulich</p> <p>Telephone No.: (41-22) 338.83.38</p>
--	---

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07K 14/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/40603</b> <b>(43) International Publication Date:</b> 13 July 2000 (13.07.00)
<b>(21) International Application Number:</b> PCT/US99/29213 <b>(22) International Filing Date:</b> 9 December 1999 (09.12.99) <b>(30) Priority Data:</b> 60/114,529 31 December 1998 (31.12.98) US <b>(71) Applicant (for all designated States except US):</b> GENWAY BIOTECH, INC. [US/US]; Welsh Commons, Suite E2, 1364 Welsh Road, N. Wales, PA 19454 (US). <b>(71)(72) Applicant and Inventor:</b> DUAN, Lingxun [US/US]; 130 Gwynedd Lea Drive, N. Wales, PA 19454 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> PRODUCTION OF RECOMBINANT MONELLIN USING METHYLOTROPHIC YEAST EXPRESSION SYSTEM  <b>(57) Abstract</b> <p>The present invention relates to a single-chain monellin-like protein which is stable and which is at least 100-fold sweet as compared to sucrose on the weight basis. The present invention also relates to a nucleic acid encoding said monellin-like protein. Preferably, the nucleic acid further comprises a promoter and a signal sequence for directing expression and secretion of the encoded monellin-like protein in the methylotrophic yeast <i>Pichia pastoris</i>. The present invention further relates to a recombinant <i>Pichia pastoris</i> cell containing the nucleic acid encoding the monellin-like protein, a process for producing the monellin-like protein from the recombinant <i>Pichia pastoris</i> and product of the process.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

**PRODUCTION OF RECOMBINANT MONELLIN USING  
METHYLOTROPHIC YEAST EXPRESSION SYSTEM**

This application claims the benefit of priority under 35 U.S.C. §119(e) to  
5 U.S. provisional application Serial No. 60/114,529 to Lingxun Duan, filed  
December 31, 1998, and entitled PRODUCTION OF RECOMBINANT  
MONELLIN USING METHYLOTROPHIC YEAST EXPRESSION SYSTEM.

**1. FIELD OF THE INVENTION**

10 The present invention relates to a single-chain monellin-like protein which is  
stable and which is at least 100-fold sweet as compared to sucrose on the weight  
basis. The present invention also relates to a nucleic acid encoding said monellin-  
like protein. Preferably, the nucleic acid further comprises a promoter and a signal  
sequence for directing expression and secretion of the encoded monellin-like  
15 protein in the methylotrophic yeast *Pichia pastoris*. The present invention further  
relates to a recombinant *Pichia pastoris* cell containing the nucleic acid encoding  
the monellin-like protein, a process for producing the monellin-like protein from  
the recombinant *Pichia pastoris* and product of the process.

**2. BACKGROUND ART**

**2.1. MONELLIN**

20 Monellin belongs to a family of intensely sweet proteins derived from  
tropical plants (Dansby, *Nature Biotechnology*, 1997, 15:419-420). Monellin is  
about 3,000-fold sweet as compared to sucrose. Other similar proteins include  
25 thaumatin, miraculin, mabinlin, pentadin and aspartame (Id.) Monellin was first  
isolated from the West African Plant *Dioscoreophyllum comminisii* (U.S. Patent  
Nos. 3,878,184 and 3,998,798; Morris and Cagan, *Biochim. Biophys. Acta*, 1972,  
261:114-122). The amino acid sequence, the three-dimensional structure and  
various physical and chemical properties of monellin have been characterized  
30 (Ogata, et al., *Nature*, 1987, 328:739-742; Morris et al., *J. Biol. Chem.*, 1973,  
248:534-539; Cagan, *Science*, 1973, 181:32-35; Bohak and Li, *Biochim. Biophys.*  
*Acta*, 1976, 427:153-170; Hudson and Beeman, *Biochem. Biophys. Res. Comm.*,  
1976, 71:212-220; Van der Wel and Loeve, *FEBS Lett.*, 1973, 29:181-183; and  
Frank and Zuber, *HoppeSeyler's Z Physiol. Chem.*, 1976, 357:585-592).

U.S. Patent No. 4,300,576 discloses smoking articles containing thaumatin or monellin. U.S. Patent No. 4,562,076 discloses chewing gum with coating of thaumatin or monellin. U.S. Patent No. 4,412,984 discloses flavor potentiated oral compositions containing thaumatin or monellin. However, despite its potential as  
5 low-calorie sweeteners, wide commercial application of monellin is hampered by concerns over its poor stability to heat and pH, lack of access to sources of supply of the plant and uncertainty in the regulatory climate for food additives (Dansby, *Nature Biotechnology*, 1997, 15:419-420).

In 1989, Sung-Hou Kim's group reported production of single-chain  
10 monellin in *E. coli* by genetic engineering (Kim et al., *Protein Eng.*, 1989, 2:571-575). The purified single-chain monellin was found to be more heat-stable and tolerant to a wide pH range, but retained the intensity of sweetness. Several aspects of this invention have been the subject of certain U.S. patents. For example, U.S. Patent No. 5,234,834 discloses constructs for expression of single-  
15 chain monellin in plant cells. U.S. Patent No. 5,487,923 discloses a sweet proteinaceous compound of the formula B-C-A, wherein B represents a peptide portion at least 90% homologous to residues 1-46 of the B chain of native monellin and modified only by conservative substitutions; C is a covalent bond or is a hydrophilic, physiologically acceptable covalent linker capable of providing a  
20 spacing length equivalent to a peptide of 1-10 amino acids selected so as to reside on the external portion of the molecule and not to disturb the native conformation; and A represents a peptide at least 90% homologous to residues 6-45 of the A chain of native monellin and modified only by conservative substitution. U.S. Patent No. 5,487,983 discloses an expression system for making the single-chain  
25 monellin disclosed in U.S. Patent No. 5,487,923. U.S. Patent No. 5,670,339 discloses DNA encoding the single-chain monellin disclosed in U.S. Patent No. 5,487,923. U.S. Patent No. 5,672,372 discloses methods for sweetening a food composition with the single-chain monellin disclosed in U.S. Patent No. 5,487,923. U.S. Patent No. 5,264,558 discloses a single-chain monellin protein that is, in a  
30 standard taste test, at least 50 times that of sucrose on a weight basis.

Recently, Kondo et al., *Nature Biotechnology*, 1997, 15:453-457 discloses heterologous expression of a single-chain monellin protein in the yeast *Candida utilis* intracellularly. It reports that monellin was produced at a high level, accounting for >50% of the soluble protein.

## 2.2. EXPRESSION OF HETEROLOGOUS PROTEINS IN *PICHIA PASTORIS*

The methylotrophic yeast *Pichia pastoris* has been used as a protein  
5 expression system. Several aspects of this expression system have been the subject  
of certain U.S. patents. For example, U.S. Patent No. 4,837,148 discloses  
autonomous replication sequences for *Pichia pastoris*. U.S. Patent No. 4,855,231  
discloses regulatory region for heterologous gene expression in *Pichia pastoris*  
cells. U.S. Patent No. 4,882,279 discloses site selective genomic modification of  
10 *Pichia pastoris*. U.S. Patent No. 4,929,555 discloses a method for making whole  
cells of *Pichia pastoris* competent for transformation. U.S. Patent No. 5,122,465  
discloses a process for generating a selectable phenotype in strains of *Pichia*  
*pastoris*. U.S. Patent No. 5,324,639 discloses production of insulin-like growth  
factor-1 in methylotrophic cells, including *Pichia pastoris* cells.

15 A number of signal sequences have been used to direct secretion of  
heterologous proteins expressed in *Pichia pastoris* cells. Examples of such signal  
sequences include, but are not limited to, the signal sequence of *Pichia pastoris*  
acid phosphatase, the signal sequence of *Aspergillus giganteus* alpha-Sarcin  
(Martinez-Ruiz et al., *Protein Expr. Purif.*, 1998, 12(3):315-22; Abdulaev et al.,  
20 *Protein Expr. Purif.*, 1997, 10(1):61-9; Kotake et al., *J. Lipid Res.*, 1996,  
37(3):599-605), the signal sequence of alpha-N-Acetylgalactosaminidase  
(alphaNAGAL, EC 3.2.1.49) (Zhu et al., *Arch. Biochem. Biophys.*, 1998,  
352(1):1-8), the signal peptide of the OmpA protein (Heim et al., *Biochim.*  
*Biophys. Acta.*, 1998, 1396(3):306-19), the signal sequence of the mouse  
25 alpha-factor signal (cCell) or the native signal sequence of pepper  
endo-beta-1,4-glucanases (Ferrarese et al., *FEBS Lett.*, 1998, 422(1):23-6), signal  
peptide of laccase isolated from the ligninolytic fungus *Trametes* (Jonsson et al.,  
*Curr. Genet.*, 1997, 32(6):425-30), signal peptide of murine lysosomal acid  
alpha-mannosidase (Merkle et al., *Biochim. Biophys. Acta.*, 1997, 1336(2):132-46),  
30 signal peptide of the porcine inhibitor of carbonic anhydrase (Wuebbens et al.,  
*Biochemistry*, 1997, 36(14):4327-36), signal sequence of *Aspergillus awamori*  
glucoamylase (Fierobe et al., *Protein Expr. Purif.*, 1997, 9(2):159-70), signal  
sequence of mouse major urinary protein (Ferrari et al., *FEBS Lett.*, 1997,  
401(1):73-7), signal sequence of pho1 (Skory et al., *Curr. Genet.*, 1996,

30(5):417-22), signal sequence of rabbit angiotensin-converting enzyme (ACE) (Sadhukhan et al., *J. Biol. Chem.*, 1996, 271(31):18310-3), prepeptide sequence of *Pichia pastoris* aspartic proteinase (Tsujikawa et al., *Yeast*, 1996, 12(6):541-53), signal sequence of *Pichia pastoris* PRC1 (Ohi et al., *Yeast*, 1996, 12(1):31-40), the  
5 signal sequence of a bacterial thermostable alpha amylase and SUC2 gene signal sequence from *Saccharomyces cerevisiae* (Paifer et al., *Yeast*, 1994, 10(11):1415-9) and the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\sigma$ -factor (Fidler et al., *J. Mol. Endocrinol.*, 1998, 21(3):327-336).

Although the methylotrophic yeast *Pichia pastoris* has been used  
10 successfully for the production of various heterologous proteins, U.S. Patent No. 5,324,639 discloses that at the present level of understanding of methylotrophic yeast expression systems, it is unpredictable whether a given gene can be expressed to an appreciable level in such yeast or whether the yeast host will tolerate the presence of the recombinant gene product in its cells. U.S. Patent No. 5,324,639  
15 further discloses that it is especially difficult to foresee if a particular protein will be secreted by the methylotrophic yeast host, and if it is, at what efficiency. For example, Vollmer et al., *J. Immunol. Methods*, 1996, 199(1):47-54, reports that when the 323 amino acid residues of the human sIL-6R are inserted into an expression/secretion vector suitable for the methylotrophic yeast *Pichia pastoris*, no  
20 detectable expression and secretion of the recombinant protein was obtained. Up to date, monellin has not been expressed and secreted using the *Pichia pastoris* expression system.

Given the great interest in the commercial application of monellin, there is a great need for a more efficient method for producing stable monellin which still  
25 retains its native sweet intensity and which simplify down stream purification procedures. The present invention addresses these and other needs in the art. Citation of references hereinabove shall not be construed as an admission that such references are prior art to the present invention.

### 30 **3. SUMMARY OF THE INVENTION**

The present invention relates to an isolated nucleic acid comprising a nucleotide sequence encoding a chimeric protein, said chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-50 of the B chain of

native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the B chain of native monellin; b) a peptidyl bond, or a second peptidyl fragment consisting of 1-12 amino acids; and c) a third peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-45 of the A chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the A chain of native monellin, wherein said chimeric protein is stable and a given amount of said chimeric protein is at least 100-fold sweet as compared to the identical amount of sucrose, and within said nucleic acid, codons which are preferably used by yeast cells are used. Preferably, the isolated nucleic acid further encodes a promoter which is capable of directing protein expression in *Pichia pastoris* and/or an amino acid sequence which is capable of directing secretion of the encoded chimeric protein from *Pichia pastoris*.

The present invention also relates to a recombinant *Pichia pastoris* cell containing the above nucleic acids. The present invention further relates to a process for producing a chimeric protein comprising growing a recombinant *Pichia pastoris* cell containing the above nucleic acid such that the encoded chimeric protein is expressed and secreted by the cell, and recovering the expressed and secreted chimeric protein. Finally, the present invention relates to products of the above processes.

#### 4. BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the amino acid sequence of a recombinant single-chain monellin protein and the DNA sequence encoding the recombinant single-chain monellin protein. Amino acid residues 1-50 corresponds to the amino acid residues 1-50 of the B chain of native monellin; amino acid residue 51 is Glycine as the linker; and amino acid residues 52-96 corresponds to the amino acid residues 1-45 of the A chain of native monellin.

FIG. 2 shows the DNA sequence of the oligos which were used for synthesis of the recombinant single-chain monellin protein.

FIG. 3 shows the location of each of DNA oligo in the synthesized monellin DNA and its enzymatic digestion sites.

FIG. 4 shows the restriction map of recombinant monellin protein expression vector pGWYS1.



FIG.5 shows the construction of pGWYS1.

FIG.6 shows the SDS-PAGE analysis of the secreted recombinant monellin protein isolated from the culture medium.

FIG.7 shows the steps for purifying the secreted recombinant monellin protein.

## 5. DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a nucleic acid encoding a single-chain monellin-like protein which is stable and which is at least 100-fold sweet as compared to sucrose on the weight basis. Preferably, the nucleic acid further comprises a promoter and a signal sequence for directing expression and secretion of the encoded monellin-like protein in the methylotrophic yeast *Pichia pastoris*. The present invention also provides a recombinant *Pichia pastoris* cell containing the nucleic acid encoding the monellin-like protein, a process for producing the monellin-like protein from the recombinant *Pichia pastoris* and product of the process.

For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the subsections which follow.

### 5.1. NUCLEIC ACIDS ENCODING THE SINGLE-CHAIN MONELLIN PROTEINS

The present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding a chimeric protein, said chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-50 of the B chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the B chain of native monellin; b) a peptidyl bond, or a second peptidyl fragment consisting of 1-12 amino acids; and c) a third peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-45 of the A chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the A chain of native monellin, wherein said chimeric protein is stable and a given amount of said chimeric protein is at least 100-fold sweet as compared to the identical amount of

sucrose, and within said nucleic acid, codons which are preferably used by yeast cells are used.

In a specific embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein wherein the  
5 first peptidyl fragment consists of an amino acid sequence that has at least 60% identity to the B chain of native monellin. Preferably, the first peptidyl fragment consists of an amino acid sequence that has at least 90% identity to the B chain of native monellin. More preferably, the first peptidyl fragment consists of the amino acid residues 1-50 of the B chain of native monellin.

10 In another specific embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein wherein the second peptidyl fragment consists of the amino acid sequence Gly-Gly-Gly-Ser-Gly-Gly-Gly-Ser (SEQ ID NO:1). Preferably, the second peptidyl fragment consists of the amino acid sequence Gly-Gly-Gly-Ser (SEQ ID  
15 NO:2). More preferably, the second peptidyl fragment consists of amino acid residue Gly.

In still another specific embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein wherein the third peptidyl fragment consists of an amino acid sequence that  
20 has at least 60% identity to the A chain of native monellin. Preferably, the third peptidyl fragment consists of an amino acid sequence that has at least 90% identity to the A chain of native monellin. More preferably, the third peptidyl fragment consists of the amino acid residues 1-45 of the A chain of native monellin.

In a preferred embodiment, the present invention provides an isolated  
25 nucleic acid comprising a nucleotide sequence encoding the chimeric protein wherein the first peptidyl fragment consists of the amino acid residues 1-50 of the B chain of native monellin, the second peptidyl fragment consists of the amino acid residue Gly and the third peptidyl fragment consists of the amino acid residues 1-45 of the A chain of native monellin.

30 In a specific embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein which is

capable of being immunoreactively bound by an anti-monellin or an anti-thaumatococcus antibody.

In another specific embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein

5 wherein the chimeric protein further comprises an amino acid sequence which is capable of directing secretion of said chimeric protein from *Pichia pastoris*. Preferably, the secretion-directing sequence is an endogenous signal sequence of *Pichia pastoris*. More preferably, the endogenous signal sequence is selected from the group consisting of the signal sequence of *Pichia pastoris* acid phosphatase,

10 *Pichia pastoris* aspartic proteinase and *Pichia pastoris* carboxypeptidase Y encoded by *Pichia pastoris* PRC1. Alternatively, the secretion-directing sequence is a yeast signal sequence, wherein said yeast is not *Pichia pastoris*. Preferably, the yeast signal sequence is a signal sequence from *Saccharomyces cerevisiae*. More preferably, the *Saccharomyces cerevisiae* signal sequence is selected from the group

15 consisting of the signal sequence of *Saccharomyces cerevisiae* SUC 2 and *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor. Most preferably, the *Saccharomyces cerevisiae* signal sequence is the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor. Examples of other secretion-directing sequences that can be used in the present invention include, but

20 are not limited to, the signal sequence of *Aspergillus giganteus* alpha-Sarcin, alpha-N-Acetylgalactosaminidase, OmpA protein, the mouse alpha-factor (cCell1), the pepper endo-beta-1,4-glucanases, the laccase isolated from the ligninolytic fungus *Trametes*, murine lysosomal acid alpha-mannosidase, the porcine inhibitor of carbonic anhydrase, *Aspergillus awamori* glucoamylase, mouse major urinary

25 protein, pho1, rabbit angiotensin-converting enzyme (ACE), and the bacterial thermostable alpha amylase.

In a specific embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein which nucleic acid is a DNA. In another specific embodiment, the present invention provides an

30 isolated nucleic acid which is hybridizable to the DNA sequence encoding the chimeric protein. In still another specific embodiment, the present invention

provides an isolated nucleic acid comprising a nucleotide sequence complementary to the nucleotide sequence encoding the chimeric protein.

In a specific embodiment, the present invention provides a DNA encoding the chimeric protein which DNA further comprises a promoter which is capable of directing protein expression in *Pichia pastoris*. Preferably, the promoter is an endogenous promoter of *Pichia pastoris*. More preferably, the endogenous promoter is the promoter of *Pichia pastoris* glyceraldehyde-3-phosphate dehydrogenase (GAP). Alternatively, although not preferred, promoters of methanol responsive genes in methylotrophic yeast can also be used. Examples of such methanol responsive promoters include, but are not limited to, the promoter for the primary alcohol oxidase gene from *Pichia pastoris* AOX1, the promoter for the secondary alcohol oxidase gene from *Pichia pastoris* AOX2, the promoter for the dihydroxyacetone synthase gene from *Pichia pastoris* (DAS), the promoter for the P40 gene from *Pichia pastoris*, the promoter for the catalase gene from *Pichia pastoris*, and the like (see U.S. Patent No. 5,324,639).

In another specific embodiment, the present invention provides a DNA encoding the chimeric protein which DNA further includes sequences allowing for its replication and selection in bacteria. In this way, large quantities of the DNA fragment can be produced by replication in bacteria.

In a preferred embodiment, the present invention provides a DNA encoding the chimeric protein, wherein within the encoded chimeric protein, the first peptidyl fragment consists of the amino acid residues 1-50 of the B chain of native monellin, the second peptidyl fragment consists of the amino acid residue Gly and the third peptidyl fragment consists of the amino acid residues 1-45 of the A chain of native monellin, and said DNA further comprises the promoter of *Pichia pastoris* GAP and the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor.

In another preferred embodiment, the present invention provides a DNA encoding the chimeric protein, wherein the codons which are preferably used by *Pichia pastoris* cells are used.

In a most preferred embodiment, the present invention provides a DNA encoding the chimeric protein wherein the DNA molecule comprises nucleotide sequence as depicted in Figure 1 or the DNA vector as depicted in Figure 4.

The nucleic acid comprising a nucleotide sequence encoding the chimeric protein disclosed herein, or any fragments, analogues or derivatives thereof, can be obtained by any method(s) known in the art. The nucleic acid may be chemically synthesized entirely. Alternatively, the nucleic acid encoding each fragment of the chimeric protein, *i.e.*, the first, second or third peptidyl fragment, may be obtained by molecular cloning or may be purified from the desired cells. The nucleic acid encoding each fragment of the chimeric protein may then be chemically or enzymatically ligated together to form the nucleic acid comprising a nucleotide sequence encoding the chimeric protein disclosed herein, or any fragments, analogues or derivatives thereof.

Any *Dioscoreophyllum comminisii* cell potentially can serve as the nucleic acid source for the isolation of the nucleic acids encoding monellin. Alternatively, the nucleic acids encoding monellin can be designed and synthesized according to the amino acid sequence of the native monellin depicted in Figure 1 (*see also* U.S. Patent No. 5,478,923).

The DNA may be obtained by standard procedures known in the art from cloned DNA (*e.g.*, a DNA "library"), by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell (*See*, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, DNA Cloning: A Practical Approach, MRL Press, Ltd., Oxford, U.K. Vol. I, II.) Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will contain only exon sequences. Whatever the source, the gene should be molecularly cloned into a suitable vector for propagation of the gene.

In the molecular cloning of the gene from cDNA, cDNA is generated from totally cellular RNA or mRNA by methods that are well known in the art. The gene may also be obtained from genomic DNA, where DNA fragments are generated (*e.g.*, using restriction enzymes or by mechanical shearing), some of

which will encode the desired gene. The linear DNA fragments can then be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once a nucleic acid comprising a nucleotide sequence encoding the chimeric protein disclosed herein, or any fragments, analogues or derivatives thereof, has been obtained, its identity can be confirmed by nucleic acid sequencing (by any method well known in the art) and comparison to the known sequences. DNA sequence analysis can be performed by any techniques known in the art, including but not limited to the method of Maxam and Gilbert (Maxam and Gilbert, 1980, *Meth. Enzymol.*, 65:499-560), the Sanger dideoxy method (Sanger et al., 1977, *Proc. Natl. Acad. Sci. U.S.A.*, 74:5463), the use of T7 DNA polymerase (Tabor and Richardson, U.S. Patent No. 4,795,699), use of an automated DNA sequenator (e.g., Applied Biosystems, Foster City, CA) or the method described in PCT Publication WO 97/15690.

Nucleic acids which are hybridizable to a nucleic acid comprising a nucleotide sequence encoding the chimeric protein disclosed herein, or any fragments, analogues or derivatives thereof, can be isolated, by nucleic acid hybridization under conditions of low, high, or moderate stringency (*See also* Shilo and Weinberg, 1981, *Proc. Natl. Acad. Sci. USA*, 78:6789-6792).

As used herein, "stable" means that a claimed single-chain monellin chimeric protein retains at least 70% of its sweet intensity after the protein has been placed at about 4°C for at least 6 months, or at about 60°C for at least 2.5 hours, or at about 100°C for at least 5 minutes. In addition, "stable" means that a claimed single-chain monellin chimeric protein retains at least 70% of its sweet intensity after the protein has been placed at a pH ranging from about 2.0 to about 11.0 for at least 6 hours.

Sweetness of the claimed single-chain monellin chimeric protein can be assessed using an ordinary taste test that is known in the art. For example, comparison to the sweetness of sucrose can be made by suitable dilutions on a weight basis (*see also* U.S. Patent No. 5,478,923).

The preferred codon usage by yeast cells can be determined by methods known in the art, e.g., methods disclosed in Sharp et al., *Nucleic Acids Res.*, 1986,

14(13):5125-43 and in Li and Luo, *J. Theor. Biol.*, 1996, 181(2):111-24.

According to Sharp, important characteristics of the preferred codon in yeast include a higher correlation with tRNA abundance, a greater degree of third base pyrimidine bias, and a lesser tendency to the A+T base pairs. Li and Luo discloses  
5 a method of classifying and predicting the gene expression level in *E. coli* and yeast cells which is called the Self-Consistent Information Clustering (SCIC).

Using the modified Codon Adaption Index (CAI) values, Li and Luo have accomplished the linear regression analysis on the relation between base composition, base correlation and gene expression level in *Escherichia coli* and  
10 yeast. Li and Luo also proposed the assumption of Expression-Enhancing-Network Site (EENS), the existence of which can be demonstrated by the linear equations between gene expression and base correlations in a codon, in adjacent codons and in non-adjacent codons. In addition, the codons that have been successfully used for expressing heterologous proteins in *Pichia pastoris* cells can be used. Examples

15 of such codons can be found in U.S. Patent No. 4,837,148; U.S. Patent No. 4,855,231; U.S. Patent No. 4,882,279; U.S. Patent No. 4,929,555; U.S. Patent No. 5,122,465; U.S. Patent No. 5,324,639; Martinez-Ruiz et al., *Protein Expr. Purif.*, 1998, 12(3):315-22; Abdulaev et al., *Protein Expr. Purif.*, 1997, 10(1):61-9; Kotake et al., *J. Lipid Res.*, 1996, 37(3):599-605; Zhu et al., *Arch. Biochem. Biophys.*, 1998, 352(1):1-8; Heim et al., *Biochim. Biophys. Acta.*, 1998, 1396(3):306-19; Ferrarese et al., *FEBS Lett.*, 1998, 422(1):23-6; Jonsson et al., *Curr. Genet.*, 1997, 32(6):425-30; Merkle et al., *Biochim. Biophys. Acta.*, 1997, 1336(2):132-46; Wuebbens et al., *Biochemistry*, 1997, 36(14):4327-36; Fierobe et al., *Protein Expr. Purif.*, 1997, 9(2):159-70; Ferrari et al., *FEBS Lett.*, 1997, 401(1):73-7; Skory et al., *Curr. Genet.*, 1996, 30(5):417-22; Sadhukhan et al., *J. Biol. Chem.*, 1996, 271(31):18310-3; Tsujikawa et al., *Yeast*, 1996, 12(6):541-53; Ohi et al., *Yeast*, 1996, 12(1):31-40; Paifer et al., *Yeast*, 1994, 10(11):1415-9; Fidler et al., *J. Mol. Endocrinol.*, 1998, 21(3):327-336; and Brocca et al., *Protein Sci.*, 1998, 7(6):1415-22.

30 Whether a chimeric protein is capable of being immunoreactively bound by an anti-monellin or an anti-thaumatococcus antibody can be determined by methods known in the art. The examples of anti-monellin or an anti-thaumatococcus antibodies

that can be used in the present invention include, but are not limited to, the antibodies disclosed in Slootstra et al., *Chem. Senses*, 1995, 20(5):535-43; Antonenko and Zanetti, *Life Sci.*, 1994, 55(15):1187-92; Bodani et al., *Hybridoma*, 1993, 12(2):177-83; Mandal et al., *Hybridoma*, 1991, 10(4):459-66 and Haimovich, *Isr. J. Med. Sci.*, 1975, 11(11):1183.

## **5.2. PRODUCTION OF MONELLIN PROTEINS FROM RECOMBINANT PICHIA PASTORIS CELLS**

In a specific embodiment, the present invention provides a recombinant *Pichia pastoris* cell containing the nucleic acid which encodes a chimeric protein, said chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-50 of the B chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the B chain of native monellin; b) a peptidyl bond, or a second peptidyl fragment consisting of 1-12 amino acids; and c) a third peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-45 of the A chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the A chain of native monellin, wherein said chimeric protein is stable and a given amount of said chimeric protein is at least 100-fold sweet as compared to the identical amount of sucrose, and within said nucleic acid, codons which are preferably used by yeast cells are used. Preferably, the recombinant *Pichia pastoris* cell contains a DNA molecule comprises nucleotide sequence as depicted in Figure 1 or a DNA vector as depicted in Figure 4. Recombinant *Pichia pastoris* cells containing the nucleic acids disclosed in Section 4.1. are also provided.

Methods for transforming methylotrophic yeast, such as *Pichia pastoris*, as well as methods applicable for culturing methylotrophic yeast cells containing in their genome a gene encoding a heterologous protein, are known generally in the art. Preferably, the transformation, positive transformant selection and culturing methods disclosed in U.S. Patent No. 4,837,148; U.S. Patent No. 4,855,231; U.S. Patent No. 4,882,279; U.S. Patent No. 4,929,555; U.S. Patent No. 5,122,465; U.S. Patent No. 5,324,639 can be used in the present invention.



In another specific embodiment, the present invention provides a process for producing a monellin chimeric protein comprising growing a recombinant *Pichia pastoris* cell containing the nucleic acid disclosed in Section 4.1. such that the encoded chimeric protein is expressed and secreted by the cell, and recovering the expressed and secreted chimeric protein. Preferably, the recombinant *Pichia pastoris* cell containing a DNA molecule comprises nucleotide sequence as depicted in Figure 1 or a DNA vector as depicted in Figure 4 is used.

Any suitable fermentation process in the art can be used in the present process. For large-scale production of recombinant DNA-based products driven by a GAP promoter in methylotrophic yeast such as *Pichia pastoris*, a three-stage, high cell-density fed batch fermentation system is preferably employed. In the first or growth stage of this fermentation system, the expression host *Pichia pastoris* cells are cultured in defined minimal medium such as BMGY (Buffered Minimal Glycerol-complex medium) with an excess of a non-inducing carbon source (e.g., glycerol). When the expression host *Pichia pastoris* cells are grown on such carbon sources, heterologous gene expression is repressed, which allows the generation of cell mass in the absence of heterologous protein expression. During this growth stage, it is also preferred that the pH of the medium be maintained at about 5. Next, the expression host *Pichia pastoris* cells are grown on limited non-inducing carbon source for a short period of time to further increase the cell mass and to depress the glucose responsive promoter. The pH of the medium during this limited growth period is kept below 4, preferably in the range from about 2.0 to about 3.5. The final stage is the production stage wherein either the "glucose excess fed-batch mode" or the "mixed-feed fed-batch mode" can be used. In the "glucose excess fed-batch mode," 2% glucose alone is added. In the "mixed-feed fed-batch mode," a limiting amount of a non-inducing carbon source and glucose is added in the fermentor to induce the expression of the monellin gene driven by a GAP promoter.

The secreted monellin chimeric proteins can be recovered from the *Pichia pastoris* culture medium by any methods known in the art. For example, methods disclosed in U.S. Patent Nos. 3,878,184 and 3,998,798; Morris and Cagan, *Biochim. Biophys. Acta*, 1972, 261:114-122; Kim et al., *Protein Eng.*, 1989, 2:571-

575; and Recently, Kondo et al., *Nature Biotechnology*, 1997, 15:453-457 can be used for recovering and isolating the secreted monellin chimeric proteins.

Preferably, the expressed and secreted chimeric protein is recovered by a means comprising ion-exchange chromatography. More preferably, the expressed and secreted chimeric protein is recovered by a means comprising CM-Sephadex  
5 column chromatography or DEAE-Sephadex column chromatography.

In another specific embodiment, the present invention provides the product of the above processes.

10

## 6. EXAMPLE

### 6.1. Preparation of the Synthetic Recombinant Monellin DNA

The amino acid sequence of the recombinant monellin protein and the nucleotide sequence of the DNA encoding the recombinant monellin protein are shown in Figure 1. As shown in Figure 1, nucleotides 1-150 encode residues 1-50  
15 of the B chain of the native monellin protein; nucleotides 150-152 encode Glycine as the linking "L" portion; and nucleotides 153-287 encode residues 1-45 of the A chain of the native monellin protein. The recombinant monellin protein is preceded by the following amino acid sequence, which corresponds to a Met residue and the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor: Met-Leu-  
20 Leu-Phe-Ile-Asn-Thr-Thr-Ile-Ala-Ser-Ile-Ala-Ala-Lys-Glu-Glu-Gly-Val-Ser-Leu-Glu-Lys-Arg-Glu-Ala-Glu-Ala-Glu-Phe (SEQ ID NO:3).

This synthetic DNA encoding the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor and the recombinant monellin protein was prepared from the oligos M1-M4 and N1-N4, which were synthesized using the  
25 Applied Biosystems 380B DNA Synthesizer by ACTG company (see Figures 2-3 and 5). The oligos were isolated by urea-polyacrylamide gel electrophoresis and purified by passing through a Sep-pak C18 column (Whatman) and annealed and ligated as shown in Figure 3 to obtain the synthetic DNA bracketed by EcoRI sites.

To synthesize the DNA encoding the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor and the recombinant monellin protein, in 100  
30 ul PCR reaction volume, 2 pM of each of the oligo M2 to N3 were mixed with 10 pM M1 and N4, heated to 94°C for 5 minutes in the absence of Tag DNA

polymerase. The reaction mixture was then slowly cooled down to 37°C. After 1 unit of the Vent DNA polymerase (New England Biolabs, Inc.) was added, the PCR reaction was performed according to the standard protocol. One hundred microliter PCR reaction mixture contains 50 mM Tris-HCl (pH 8.0), 2.5 mM MgCl<sub>2</sub>, 10 mM DTT, 1 mM dNTP, and 1 unit of the Vent DNA polymerase. The PCR reaction was performed as following: at 94°C for 1 minute, 53°C for 1.5 minutes, 72°C for 2 minutes within each cycle; and for a total of 30 cycles. Finally, the reaction mixture was incubated at 72°C for 10 minutes. The reaction mixture was extracted by phenol/chloroform, precipitated with ethanol, and gel purified in 1.2% low-melting agarose gel. The purified DNA fragment was inserted into the pT7bleu (R) vector (Novagen, Inc.) to generate the pT7yM plasmid (*see* Figure 5). In 20 ul DNA ligation reaction, 2 ul of 10 mM ATP, 40 units of the T4 DNA ligase (New England Biolab, Inc.) was added and mixed with 1 ug purified monellin DNA fragment and 50 ng pT7blue (R) vector. The reaction mixture was kept at 16°C for 16 hours. The ligation mixture was transformed into host cells by adding 5 ul of the ligation mixture to 200 ul of *E. coli* NovaBlue competent cells (Messing, *Methods in Enzymology*, 1983, 101:20-78) and the desired sequence was confirmed by dideoxy sequencing using T7 and U19 primer (Sanger et al., *Proc. Natl. Acad. Sci.*, 1985, 74:5463-5467).

(1)

## **6.2. Preparation of the Expression Vector pGWYS-1**

The pGAPZa expression vector was purchased from Invitrogen, Inc. The synthetic monellin DNA fragment was removed from pT7yMenallin with EcoRI and inserted into an EcoRI site of the pGAPZa vector to give pGWYS. Briefly, 5 ug purified pT7yMenallion plasmid was digested in 20 ul reaction volume using 5 units EcoRI (Promega Inc.) at 37°C for 2 hours. After the reaction mixture was separated by 1% low-melting agarose gel electrophoresis, the synthetic monellin DNA fragment was purified using the Wizard PCR Preps DNA purification kit (Promega, Inc). One hundred ng purified monellin DNA fragment were used for ligation into the expression pGAPZa vector. In the 10 ul ligation reaction, 50 ng of the EcoRI digested pGAPZa vector was mixed with 100 ng purified monellin DNA fragment. The ligation reaction was carried out in the presence of 10 ul of

20 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 10 mM DTT and 200 units of the T4 DNA ligase (New England Biolabs, Inc.) at 16°C overnight to give the pGWYS. The ligation mixture was transformed into *E. coli* TOP10F' cells (Invitrogen, Inc.). Twenty Zeocin-resistant clones were picked and orientation of the insert was  
5 screened by PCR reaction using the  $\alpha$ -factor ( 5'CTATTGCCAGCATTGCTGC3') (SEQ ID NO:4) and the N4 oligos. Each selected clone was transferred into 3 ml LB culture medium containing 200 ug/ml Zeocin and incubated with shaking at 37°C overnight. The recombinant plasmid pGWYS was prepared from 1.5 ml  
10 cultured bacteria cells using the Qiagen Tip 20 kit system (Qiagen, Inc.). Fifty ng purified pGWYS plasmid was used as the PCR template to determine orientation of the insert. In 25 ul PCR reaction, 50 ng of the pGWYS plasmid was mixed with 2.5 pM of the  $\alpha$ -factor and the N4 oligos in the presence of 1 unit Taq DNA polymerase (Promega, Inc.). The PCR reaction was performed under the following conditions: at 94°C for 1 minute, 55°C. for 1 minute, 72°C for 2 minutes within  
15 each cycle; and for a total of 40 cycles. Finally, the reaction mixture was incubated at 72°C. for 10 minutes. After the PCR reaction mixture was separated on 1.2% agarose gel, one of the clones which contains the insert with the desired orientation was named pGWYS-1. The sequence of the insert was further confirmed by DNA sequencing.

20

### **6.3. Transformation of *Pichia pastoris* Cells with the pGWYS-1**

To generate the high-level and stable expression of monellin in *Pichia pastoris*, purified pGWYS-1 plasmid was transformed into *Pichia pastoris* cells by electroporation technique described in the *Pichia pastoris* Expression Kit Manual  
25 using the Electroporation Apparatus II (Invitrogen Inc.). Briefly, 500 ml of the *Pichia pastoris* GS115 cells were grown in YPD medium at 30°C to an OD<sub>600</sub> of 1.3. Cells were pelleted with a centrifugation of 1,500 g for 5 minutes at 4°C. Pelleted cells were washed with 500ml of ice-cold sterile water. The washing step was repeated with 250 ml and 20 ml ice-cold sterile water, receptively. Then, the  
30 cells were washed with 20 ml of ice-cold 1 M sorbitol and resuspend in 1 ml ice-cold sorbitol. Forty ul of the yeast GS115 cells in 1 M sorbitol were mixed with 10 ug purified pGWYS-1 plasmid to total volume 50 ul and the mixture was

transferred into an ice-cold cuvette. The cuvette containing the mixture was incubated on ice for 5 minutes. Electroporation was performed according to the Electroporation Apparatus II manual parameters (Invitrogen Inc manufacture). After the electrical pulse, 1 ml of ice-cold 1M sorbitol was added into the cuvette, and the content of the cuvette was transferred into a microcentrifuge tube. Two hundred  $\mu$ l transformed cells were plated on one 5 RDB plate containing 400  $\mu$ g/ml Zeocin. The plates were incubated at 30°C until colonies appeared. Positive transformants were characterized by their growth in the presence of Zeocin at various concentrations, *e.g.*, 400  $\mu$ g/ml, 600  $\mu$ g/ml, 800  $\mu$ g/ml and 1000 $\mu$ g/ml.

#### 6.4. Stability Test of the Positive Transformants

Three positive transformants were selected for further characterization based on their growth in the presence of 800 $\mu$ g/ml Zeocin and the expression of recombinant monellin by the 2% glucose induction. The following experiment was performed to test their genetic stability. Each of these 3 positive transformants was picked up using a sterile toothpick and incubated on a YPD plate without any selection at 30°C until colonies appeared. The colonies were picked up and plated on a new YPD plate until new colonies appeared. After such non-selective growth was repeated 50 times, each of the passage colonies was incubated on a selective plate containing 800  $\mu$ g/ml Zeocin. The protein expression upon 2% glucose induction was analyzed by SDS-PAGE. All three positive transformants showed the same phenotype as the original colonies after 50 times passage on the YPD plates.

#### 6.5 Production of the Recombinant Monellin Protein

Each of the three positive transformants selected in 5.4. was grown in 1 liter YPD medium in 5 liter flask at 30°C with vigorous shaking (250rpm). Two ml supernatant were obtained from the culture after 24 hours, 48 hours and 72 hours, respectively. Five  $\mu$ l of the samples collected at each time point were analyzed using the 15-20% gradient polyacrylamide gel. The secreted recombinant monellin protein was observed as the 12 kD protein band. Quantitation of the SDS-PAGE analysis using the Densitometer (Molecular Dynamic, Inc.) indicates that one of the

positive strain produced nearly 10 grams per liter secreted recombinant single-chain monellin protein. This strain was named GWyS1.

#### 6.6 Purification of Secreted Recombinant Monellin Protein

5 Protein methods were used to purify the recombinant monellin protein secreted from GwyS-1 yeast strain. According to the first method, after 72-hour culturing, supernatant was collected by a centrifugation at 12,000 rpm (17,000g). After collection, the supernatant pH was adjusted to about 6.8 using 0.1 N NaOH solution. One M  $\text{NaH}_2\text{PO}_4$ - $\text{Na}_2\text{HPO}_4$  (pH6.8) was added into the supernatant till  
10 1:100 (v/v) and mixed well. The supernatant was then loaded on the CM-Sephadex column (Pharmacia, Inc.) pre-equilibrated with 0.01 M  $\text{NaH}_2\text{PO}_4$ - $\text{Na}_2\text{HPO}_4$  (pH6.8) solution. After the column was washed with 5 column volume 0.01 M  $\text{NaH}_2\text{PO}_4$ - $\text{Na}_2\text{HPO}_4$  (pH6.8) solution, the recombinant monellin protein was eluted with 0.3 M NaCl-0.01 M  $\text{NaH}_2\text{PO}_4$ - $\text{Na}_2\text{HPO}_4$  (pH6.8) solution. After dialysis against water, the  
15 purity of the protein was determined to be about 98% by gel electrophoresis.

According to the second method, after 72-hour culturing, supernatant was collected by a centrifugation at 12,000 rpm (17,000g). After collection, the supernatant pH was adjusted to about 7.2 using 0.1 N NaOH solution. One M NaCl-1 M  $\text{NaH}_2\text{PO}_4$ - $\text{Na}_2\text{HPO}_4$  (pH 7.2) was added into the supernatant till 1:100  
20 (v/v) and mixed well. The supernatant was then loaded on the DEAE-Sephadex column (Pharmacia, Inc.) pre-equilibrated with 1 M  $\text{NaH}_2\text{PO}_4$ - $\text{Na}_2\text{HPO}_4$  (pH 7.2)-1M NaCl solution. The flow-through fraction was collected and dialyzed against water. The purity of the protein was determined to be about 98% by gel electrophoresis.

The recombinant monellin protein purified according to either method was  
25 further lyophilized to dry powder for testing its sweetness.

#### 6.7. Sweetness and Stability Test

Sweetness of the purified recombinant monellin protein was assessed using an ordinary taste test. Comparison to the sweetness of sucrose was made by  
30 suitable dilutions on a weight basis. In a typical test, 1, 10, 25 and 50 mg/ml aqueous sucrose solutions were used as standard solutions. The minimum weight of the purified recombinant monellin protein which could generate sweet taste was compared with that of sucrose. The recombinant monellin of this invention

requires the addition of an amount which is about 1000-fold less than that of sucrose. For example, 50 ng/ml recombinant monellin protein solution was as sweet as 50 mg/ml sucrose (Lucky Supermarket's Lady Lee brand sugar).

Stability was measured by dissolving natural monellin (Sigma, Inc.) and the  
5 purified recombinant monellin protein at 100 ug/ml concentration at pH 2.0, 4.0, 6.3. and 7.5. Each sample was heated to 37°C, 50°C, 60°C, 70°C, 80°C, 90°C and 100°C for 15 minutes and let cool to room temperature before tasting. The most dramatic difference was that natural monellin lost its sweetness when heated to 50°C at pH 2.0, while the purified recombinant monellin protein retained its  
10 sweetness even after heating at 100°C for 5 minutes.

The present invention is not to be limited in scope by the microorganism deposited or the specific embodiments described herein. Indeed, various  
15 modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

Various references are cited herein, the disclosures of which are  
20 incorporated by reference in their entireties.

**WHAT IS CLAIMED IS:**

1. An isolated nucleic acid comprising a nucleotide sequence encoding a chimeric protein, said chimeric protein comprises, from N-terminus to C-terminus:
- a first peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-50 of the B chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the B chain of native monellin;
  - a peptidyl bond, or a second peptidyl fragment consisting of 1-12 amino acids; and
  - a third peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-45 of the A chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the A chain of native monellin,
- wherein said chimeric protein is stable and a given amount of said chimeric protein is at least 100-fold sweet as compared to the identical amount of sucrose, and within said nucleic acid, codons which are preferably used by yeast cells are used.
2. The isolated nucleic acid of claim 1, wherein the first peptidyl fragment consists of an amino acid sequence that has at least 60% identity to the B chain of native monellin.
3. The isolated nucleic acid of claim 1, wherein the first peptidyl fragment consists of an amino acid sequence that has at least 90% identity to the B chain of native monellin.
4. The isolated nucleic acid of claim 1, wherein the first peptidyl fragment consists of the amino acid residues 1-50 of the B chain of native monellin depicted as the amino acid residues 1-50 in Figure 1 (SEQ ID NO:5).
5. The isolated nucleic acid of claim 1, wherein the second peptidyl fragment consists of the amino acid residue Gly.



6. The isolated nucleic acid of claim 1, wherein the second peptidyl fragment consists of the amino acid sequence Gly-Gly-Gly-Ser (SEQ ID-NO:2).

7. The isolated nucleic acid of claim 1, wherein the second  
5 peptidyl fragment consists of the amino acid sequence Gly-Gly-Gly-Ser-Gly-Gly-Gly-Ser-Gly-Gly-Ser (SEQ ID NO:1).

8. The isolated nucleic acid of claim 1, wherein the third peptidyl fragment consists of an amino acid sequence that has at least 60% identity to the A chain of native monellin.

10 9. The isolated nucleic acid of claim 1, wherein the third peptidyl fragment consists of an amino acid sequence that has at least 90% identity to the A chain of native monellin.

10. The isolated nucleic acid of claim 1, wherein the third peptidyl fragment consists of the amino acid residues 1-45 of the A chain of native  
15 monellin depicted as the amino acid residues 52-96 in Figure 1 (SEQ ID NO:5).

11. The isolated nucleic acid of claim 1 which nucleic acid encodes the amino acid residues 1-96 of Figure 1 (SEQ ID NO:5).

12. The isolated nucleic acid of claim 1, wherein the chimeric protein is capable of being immunoreactively bound by an anti-monellin antibody.

20 13. The isolated nucleic acid of claim 1, wherein the chimeric protein is capable of being immunoreactively bound by an anti-thaumatococcus antibody.

14. The isolated nucleic acid of claim 1, wherein the chimeric protein further comprises an amino acid sequence which is capable of directing secretion of said chimeric protein from *Pichia pastoris*.

25 15. The isolated nucleic acid of claim 14, wherein the secretion-directing sequence is an endogenous signal sequence of *Pichia pastoris*.

16. The isolated nucleic acid of claim 15, wherein the endogenous signal sequence is selected from the group consisting of the signal sequence of *Pichia pastoris* acid phosphatase, *Pichia pastoris* aspartic proteinase  
30 and *Pichia pastoris* carboxypeptidase Y encoded by *Pichia pastoris* PRC1.

17. The isolated nucleic acid of claim 14, wherein the secretion-directing sequence is a yeast signal sequence, wherein said yeast is not *Pichia pastoris*.

18. The isolated nucleic acid of claim 17, wherein the yeast  
5 signal sequence is a signal sequence from *Saccharomyces cerevisiae*.

19. The isolated nucleic acid of claim 18, wherein the *Saccharomyces cerevisiae* signal sequence is selected from the group consisting of the signal sequence of *Saccharomyces cerevisiae* SUC 2 and *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor.

10 20. The isolated nucleic acid of claim 19, wherein the *Saccharomyces cerevisiae* signal sequence is the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor.

21. The isolated nucleic acid of claim 11, further comprising an amino acid sequence which is capable of directing secretion of said chimeric  
15 protein from *Pichia pastoris*.

22. The isolated nucleic acid of claim 21, wherein the secretion-directing sequence is the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor.

23. The isolated nucleic acid of claim 14, wherein the secretion-  
20 directing sequence is selected from the group consisting of the signal sequence of *Aspergillus giganteus* alpha-Sarcin, alpha-N-Acetylgalactosaminidase, OmpA protein, the mouse alpha-factor (cCell1), the pepper endo-beta-1,4-glucanases, the laccase isolated from the ligninolytic fungus *Trametes*, murine lysosomal acid alpha-mannosidase, the porcine inhibitor of carbonic anhydrase, *Aspergillus*  
25 *awamori* glucoamylase, mouse major urinary protein, pho1, rabbit angiotensin-converting enzyme (ACE), and the bacterial thermostable alpha amylase.

24. The nucleic acid of claim 1, wherein said nucleic acid is a DNA.

30 25. An isolated nucleic acid comprising a nucleotide sequence complementary to the nucleotide sequence of claim 1.

26. An isolated nucleic acid hybridizable to the DNA sequence of claim 24.

27. The DNA of claim 24, further comprising a promoter which is capable of directing protein expression in *Pichia pastoris*.

5 28. The DNA of claim 27, wherein the promoter is an endogenous promoter of *Pichia pastoris*.

29. The DNA of claim 28, wherein the endogenous promoter is the promoter of *Pichia pastoris* glyceraldehyde-3-phosphate dehydrogenase.

30. The DNA of claim 24 said DNA encodes the amino acid  
10 residues 1-96 of Figure 1 (SEQ ID NO:5) and said DNA further comprises the promoter of *Pichia pastoris* glyceraldehyde-3-phosphate dehydrogenase and the signal sequence of *Saccharomyces cerevisiae* mating pheromone  $\alpha$ -factor.

31. The DNA of claim 24, wherein the codons which are preferably used by *Pichia pastoris* cells are used.

15 32. A DNA molecule comprises nucleotide sequence as depicted in Figure 1.

33. The pGWYS1 DNA vector as depicted in Figure 4.

34. A recombinant *Pichia pastoris* cell containing the nucleic acid of claim 1.

20 35. A recombinant *Pichia pastoris* cell containing the DNA of claim 32.

36. A recombinant *Pichia pastoris* cell containing the DNA of claim 33.

37. A process for producing a chimeric protein comprising  
25 growing a recombinant *Pichia pastoris* cell containing the nucleic acid of claim 1 such that the encoded chimeric protein is expressed and secreted by the cell, and recovering the expressed and secreted chimeric protein.

38. A process for producing a chimeric protein comprising  
30 growing a recombinant *Pichia pastoris* cell containing the DNA of claim 32 such that the encoded chimeric protein is expressed and secreted by the cell, and recovering the expressed and secreted chimeric protein.

39. A process for producing a chimeric protein comprising growing a recombinant *Pichia pastoris* cell containing the DNA of claim 33 such that the encoded chimeric protein is expressed and secreted by the cell, and recovering the expressed and secreted chimeric protein.

5 40. The process of claim 37, wherein the expressed and secreted chimeric protein is recovered by a means comprising ion-exchange chromatography.

41. The process of claim 40, wherein the ion-exchange chromatography being used is CM-Sephadex column chromatography.

10 42. The process of claim 40, wherein the ion-exchange chromatography being used is DEAE-Sephadex column chromatography.

43. The product of the process of claim 37.

44. The product of the process of claim 38.

45. The product of the process of claim 39.

15 46. The product of the process of claim 40.

47. The product of the process of claim 41.

48. The product of the process of claim 42.

49. A chimeric protein, said chimeric protein comprises, from N-terminus to C-terminus:

20 a) a first peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-50 of the B chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the B chain of native monellin;

25 b) a peptidyl bond, or a second peptidyl fragment consisting of 1-12 amino acids; and

c) a third peptidyl fragment consisting of an amino acid sequence that has at least 40% identity to residues 1-45 of the A chain of native monellin, in which the percentage identity is determined over an amino acid sequence of identical size to the A chain of native monellin,

30 wherein said chimeric protein is stable and a given amount of said chimeric protein is at least 100-fold sweet as compared to the identical amount of sucrose, and within said nucleic acid, codons which are preferably used by yeast cells are used.

1 / 8

GGT GAG TGG GAG ATT ATT GAC ATT GGT CCA TTC ACT  
Gly Gly Trp Glu Ile Ile Asp Ile Gly Pro Phe Thr

CAA AAC TTG GGT AAG TTC GCT GTT GAC GAG GAG AAC  
Gln Asn Leu Gly Lys Phe Ala Val Asp Glu Glu Asn

AAG ATT GGT CAA TAC GGT AGA TTG ACT TTC AAC AAG  
Lys Ile Gly Gln Tyr Gly Arg Leu Thr Phe Asn Lys

GTT ATT AGA CCA TGT ATG AAG AAG ACT ATT TAC GAG  
Val Ile Arg Pro Cys Met Lys Lys Thr Ile Tyr Glu

AAC GAG GGT TCT AGA GAG ATT AAG GGT TAC GAG TAC  
Asn Glu Gly Ser Arg Glu Ile Lys Gly Tyr Glu Tyr

CAA TTG TAC GTT TAC GCT TCT GAC AAG TTG TTC CGT  
Gln Leu Tyr Val Tyr Ala Ser Asp Lys Leu Phe Arg

GCT GAC ATT TCT GAG GAC TAC AAG ACT CGT GGT CGT  
Ala Asp Ile Ser Glu Asp Tyr Lys Thr Arg Gly Arg

AAG TTG TTG AGA TTC AAC GGT CCA GTT CCA CCA CCA  
Lys Leu Leu Arg Phe Asn Gly Pro Val Pro Pro Pro

TAA  
Stop

FIG.1

2 / 8

M1

5' AGA ATT CGG TGA GTG GGA GAT TAT TGA CAT TGG TCC ATT  
CAC TCA AAA CTT GG 3'

M2

5' GAA CAA GAT TGG TCA ATA CGG TAG ATT GAC TTT CAA CAA  
GTT TAT TAG GCC ATG T 3'

M3

5' GAG ACC GAG GGT TCT AGA GAG ATT AAG GGT TAC GAG TAC  
CAA TTG TAC GTT TAC GCT TC 3'

M4

5' GTG CTG ACA TTC CTG AGG ACT ACA AGA CTC GTG GTC GTA  
AGT TGT TGA GAT TC 3'

N1

5' GTA TTG ACC AAT CTT GTT CTC CTC GTC AAC AGC GAA CTT  
ACC CAA GTT TTG AGT GAA TG 3'

N2

5' CTC TAG AAC CCT CGT TCT CGT AAA TAG TCT TCT TCA TAC  
ATG GTC TAA TAA CCT TG 3'

N3

5' GTC CTC AGA AAT GTC AGC ACG GAA CAA CTT GTC AGA AGC  
GTA AAC GTA CAA TTG

N4

5' AGA ATT CTT ATG GTG GTG GAA CTG GAC CGT TGA ATC TCA  
ACA ACT TAC GAC 3'

**FIG. 2**

3 / 8





FIG. 3

4 / 8

**Comments for pGWYS1 (3479 bp)**

GAP Promoter region: 1 -483  
 Alpha-factor signal sequence: 493-760  
 Monellin coding region: 762-1059  
 3' AOX 1 termination region: 1060-1306  
 TEF1 Promoter region: 1307-1709  
 EM7 Promoter region: 1710-1781  
 Sh ble ORF: 1782-2518  
 CYC1 termination region: 2159 2477  
 ColE1 origin ( pUC-derived): 2478-3479

**FIG. 4**



5 / 8





FIG. 5

6 / 8



Lane 1. Protein MW Marker

Lane 2. 5ul Culture Medium

Lane 3. Partially Purified Recombinant  
Single Chain Monellin

Lane 4. 40ug Native Monellin

**FIG.6**

7 / 8




FIG. 7a

8 / 8




FIG. 7b

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> C12N 15/29, 15/81, 1/19, 15/62, C07K 14/43 // (C12N 1/19, C12R 1:84)	<b>A3</b>	<b>(11) International Publication Number:</b> WO 00/40603 <b>(43) International Publication Date:</b> 13 July 2000 (13.07.00)
<b>(21) International Application Number:</b> PCT/US99/29213 <b>(22) International Filing Date:</b> 9 December 1999 (09.12.99)  <b>(30) Priority Data:</b> 60/114,529 31 December 1998 (31.12.98) US  <b>(71) Applicant (for all designated States except US):</b> GENWAY BIOTECH, INC. [US/US]; Welsh Commons, Suite E2, 1364 Welsh Road, N. Wales, PA 19454 (US).  <b>(71)(72) Applicant and Inventor:</b> DUAN, Lingxun [US/US]; 130 Gwynedd Lea Drive, N. Wales, PA 19454 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 21 September 2000 (21.09.00)
<b>(54) Title:</b> PRODUCTION OF RECOMBINANT MONELLIN USING METHYLOTROPHIC YEAST EXPRESSION SYSTEM  <b>(57) Abstract</b>  The present invention relates to a single-chain monellin-like protein which is stable and which is at least 100-fold sweet as compared to sucrose on the weight basis. The present invention also relates to a nucleic acid encoding said monellin-like protein. Preferably, the nucleic acid further comprises a promoter and a signal sequence for directing expression and secretion of the encoded monellin-like protein in the methylotrophic yeast <i>Pichia pastoris</i> . The present invention further relates to a recombinant <i>Pichia pastoris</i> cell containing the nucleic acid encoding the monellin-like protein, a process for producing the monellin-like protein from the recombinant <i>Pichia pastoris</i> and product of the process.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/29213

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/29 C12N15/81 C12N1/19 C12N15/62 C07K14/43  
 //(C12N1/19,C12R1:84)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KONDO K ET AL: "High-level expression of a sweet protein, monellin, in the food yeast <i>Candida utilis</i> " NATURE BIOTECHNOLOGY, vol. 15, no. 5, May 1997 (1997-05), pages 453-457, XP000910200 cited in the application page 454, line 12 - line 15; figure 2B	1-5, 8-10,12, 13, 24-26,49
Y		14-23, 27-31, 34-48
Y	VEDVICK TS: "Gene expression in yeast: <i>Pichia pastoris</i> " CURR. OPIN. BIOTECHNOL., vol. 2, no. 5, October 1991 (1991-10), pages 742-745, XP000915828 the whole document	14-23, 27-31, 34-48
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

19 June 2000

Date of mailing of the international search report

06/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Nichogiannopoulou, A

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/29213

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>CREGG ET AL: "Recent advances in the expression of foreign genes in Pichia pastoris"            BIO/TECHNOLOGY,            vol. 11, 1993, pages 905-910, XP002100307            the whole document</p> <p>---</p>	<p>14-23,            27-31,            34-48</p>
Y	<p>US 3 998 798 A (UNIVERSITY OF PENNSYLVANIA)            21 December 1976 (1976-12-21)            cited in the application            column 3, paragraph 2 - paragraph 3</p> <p>---</p>	<p>40-42</p>
X	<p>WO 88 10265 A (UNIVERSITY OF CALIFORNIA)            29 December 1988 (1988-12-29)            cited in the application</p> <p>Sequence 96,9% identical to SEQ ID NO:5            page 3, line 1 -page 4, line 2; claims            10,11; figure 3</p> <p>---</p>	<p>1-4,            8-10,12,            13,            24-26,49</p>
X	<p>EP 0 821 055 A (INT FLAVORS &amp; FRAGRANCES            INC) 28 January 1998 (1998-01-28)</p> <p>page 20, line 1 - line 13; figure 3B</p> <p>-----</p>	<p>1-5,            8-10,12,            13,            24-26,49</p>



# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 99/29213

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3998798	A	21-12-1976	NONE	
WO 8810265	A	29-12-1988	AT 108185 T	15-07-1994
			AU 1939788 A	19-01-1989
			BR 8807102 A	17-10-1989
			CA 1340081 A	13-10-1998
			DE 3850574 D	11-08-1994
			DE 3850574 T	27-10-1994
			EP 0323489 A	12-07-1989
			EP 0374157 A	27-06-1990
			ES 2012530 A	01-04-1990
			GR 88100384 A, B	08-03-1989
			IN 168525 A	20-04-1991
			JP 2500717 T	15-03-1990
			JP 8022874 B	06-03-1996
			JP 2504028 T	22-11-1990
			WO 8810303 A	29-12-1988
			KR 9108642 B	19-10-1991
			KR 9612512 B	20-09-1996
			US 5487983 A	30-01-1996
			US 5478923 A	26-12-1995
			US 5672372 A	30-09-1997
			US 5739409 A	14-04-1998
			US 5264558 A	23-11-1993
			US 5234834 A	10-08-1993
			US 5670339 A	23-09-1997
			US 5882884 A	16-03-1999
			AU 1994088 A	19-01-1989
			EP 0318580 A	07-06-1989
			JP 2500641 T	08-03-1990
			WO 8810271 A	29-12-1988
EP 0821055	A	28-01-1998	US 6001410 A	14-12-1999


REC'D 02 APR 2001

WIPO PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

14

Applicant's or agent's file reference n.80072		<b>FOR FURTHER ACTION</b>	See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)
International application No. PCT/US99/29213	International filing date (day/month/year) 09/12/1999	Priority date (day/month/year) 31/12/1998	
International Patent Classification (IPC) or national classification and IPC C07K14/00			
Applicant GENWAY BIOTECH, INC. et al.			
<p>1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.</p> <p>2. This REPORT consists of a total of 7 sheets, including this cover sheet.</p> <p><input type="checkbox"/> This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).</p> <p>These annexes consist of a total of sheets.</p>			
<p>3. This report contains indications relating to the following items:</p> <ul style="list-style-type: none"> <li>I <input checked="" type="checkbox"/> Basis of the report</li> <li>II <input type="checkbox"/> Priority</li> <li>III <input type="checkbox"/> Non-establishment of opinion with regard to novelty, inventive step and industrial applicability</li> <li>IV <input type="checkbox"/> Lack of unity of invention</li> <li>V <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement</li> <li>VI <input type="checkbox"/> Certain documents cited</li> <li>VII <input checked="" type="checkbox"/> Certain defects in the international application</li> <li>VIII <input checked="" type="checkbox"/> Certain observations on the international application</li> </ul>			
Date of submission of the demand  14/07/2000		Date of completion of this report  29.03.2001	
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465		Authorized officer  Nichogiannopoulou, A  Telephone No. +49 89 2399 8054	



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US99/29213

## I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).):*

### Description, pages:

1-20 as originally filed

### Claims, No.:

1-49 as originally filed

### Drawings, sheets:

1/8-8/8 as originally filed

### Sequence listing part of the description, pages:

1-5, as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US99/29213

- ☐ the description,      pages:  
☐ the claims,      Nos.:  
☐ the drawings,      sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes:	Claims	6, 7, 11, 14-23, 27-48
	No:	Claims	1-5, 8-10, 12, 13, 24-26, 49
Inventive step (IS)	Yes:	Claims	6, 7, 11, 32, 33
	No:	Claims	1-5, 8-10, 12-31, 34-49
Industrial applicability (IA)	Yes:	Claims	1-49
	No:	Claims	

2. Citations and explanations  
**see separate sheet**

## VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:  
**see separate sheet**

## VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:  
**see separate sheet**

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/US99/29213

**Re Item I**

**Basis of the report**

1. This report was established based on the application documents and the written sequence listing pages 1-5.

**Re Item V**

**Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

1. Reference is made to the following documents:
  - D1: KONDO K ET AL: 'High-level expression of a sweet protein, monellin, in the food yeast *Candida utilis*' NATURE BIOTECHNOLOGY, vol. 15, no. 5, May 1997 (1997-05), pages 453-457, XP000910200 cited in the application
  - D2: VEDVICK TS: 'Gene expression in yeast: *Pichia pastoris*' CURR. OPIN. BIOTECHNOL., vol. 2, no. 5, October 1991 (1991-10), pages 742-745, XP000915828
  - D3: CREGG ET AL: 'Recent advances in the expression of foreign genes in *Pichia pastoris*' BIO/TECHNOLOGY, vol. 11, 1993, pages 905-910, XP002100307
  - D4: US-A-3 998 798 (UNIVERSITY OF PENNSYLVANIA) 21 December 1976 (1976-12-21) cited in the application
  - D5: WO 88 10265 A (UNIVERSITY OF CALIFORNIA) 29 December 1988 (1988-12-29) cited in the application
  - D6: EP-A-0 821 055 (INT FLAVORS & FRAGRANCES INC) 28 January 1998 (1998-01-28)
2. **Novelty** (Article 33(2) PCT)

The present application discloses the production of recombinant single-chain chimeric monellin in the methylotropic yeast *Pichia pastoris*.

- 2.1. **D1** (cited in the application) discloses the heterologous expression of a single-chain monellin gene in the yeast *Candida utilis* under the control of a *C. utilis* promoter. The single-chain monellin (Fig. 2B) is of the formula B-C-A, wherein C is a linker consisting of Gly. **D1** is thus detrimental to the novelty and inventive step of claims 1-5, 8-10, 12, 13, 24-26 and 49.
- 2.2. **D5** (cited in the application) discloses single chain monellin analogues and their recombinant production. The analogues have the formula B-C-A. C is a covalent linker 3-10 amino acids long. Limited modifications of the A and B chain portions can be included. The compound with a reported sweetness at least 50 times that of sucrose has 96.9% identity to SEQ ID No:5. **D5** is thus detrimental to the novelty and inventive step of claims 1-4, 8-10, 12, 13, 24-26 and 49.
- 2.3. For the sake of completion, it is noted that **D6** also discloses single chain monellin analogues of the formula B-C-A. C is a linker that can be Gly as in the present application (see Fig. 3-B(i) and (ii)). Sequence Id 4 has 96.9% identity with the sequence of the present application while Sequence Id 7 has 97.9% identity. **D6** is thus detrimental to the novelty and inventive step of claims 1-4, 8-10, 12, 13, 24-26 and 49.
3. **Inventive step** (Article 33(3) PCT)
  - 3.1. **D2** and **D3** disclose the versatility and utility of the *Pichia pastoris* expression system. Together with the teachings of **D1** on recombinant expression of the single-chain monellin, they render the subject-matter of claims 14-23, 27-31 and 34-48 obvious.
  - 3.2. **D4** (cited in the application) discloses the purification of monellin from the crude fresh fruit of *D. cumminsii* by DEAE-cellulose and CM-cellulose ion exchange

chromatography. This teaching deprives inventive step from claims 40-42.

**4. Industrial applicability (Article 33(4) PCT)**

The subject-matter of present claims 1-49 appears to be industrially applicable under the terms of Article 33(4) PCT.

**Re Item VII**

**Certain defects in the international application**

1. Contrary to the requirements of Rule 5.1(ii) PCT, documents **D2**, **D3** and **D6** are not identified in the description and the relevant background art disclosed therein is not briefly discussed.
2. Claim 32 relies on a reference to a figure, contravening Rule 6.2(a) PCT.

**Re Item VIII**

**Certain observations on the international application**

1. Claim 1 is related to a nucleic acid wherein "codons which are preferably used by yeast cells are used". This definition is not a true technical feature of the claimed compound and renders said claim not-enabled and unclear under the terms of Articles 5 and 6 PCT respectively.
2. Claim 14 does not meet the requirements of Article 6 PCT in that the matter for which protection is sought is not clearly defined. The claim attempts to define the subject-matter in terms of the result to be achieved (secretion of said protein from *Pichia pastoris*) which merely amounts to a statement of the underlying problem.

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/US99/29213

The technical features necessary for achieving this result should be added.

3. Claim 26 relates to a nucleic acid "hybridisable to the DNA sequence of claim 24". Such a definition is too vague and lacks clarity in the sense of Article 6 PCT. It is furthermore noted that such a broad term jeopardises the novelty of the claim.